

The Airwave Health Monitoring Study

Vergnaud AC, Singh D, Spear J, Heard A, Elliott P

Department of Epidemiology & Biostatistics, Imperial College London, UK, in collaboration with Electrocardiology Core Lab, University of Glasgow; and Queen Mary, University of London

Contact e-mail: a.vergnaud@imperial.ac.uk d.singh@imperial.ac.uk

ACKNOWLEDGMENTS This study is funded by the National Policing Improvement Agency (NPIA) and the National Institute for Health Research comprehensive Biomedical Research Centre, Imperial College Healthcare NHS Trust

Objectives

The Airwave Health Monitoring Study is a large occupational prospective cohort study of the British police forces which was established to investigate possible long-term health effects associated with use of the Airwave (TETRA) radio system.

The study involves health screening with collection and storage of data and samples linked to health records with consent for their use for future research.




Methods

Population and design

- Participants are recruited among the participating police forces and agencies in England, Wales and Scotland with a target sample size of 70,000.
- Participants are invited to:
 - complete an enrolment questionnaire (22.0% response), providing data on socio-demographic characteristics, lifestyle, health and Airwave usage.
 - attend a health screening (45.7 % response), providing data on lifestyle, health and well-being, medical history, Airwave usage, 7-day diet questionnaire, cognitive test, clinical measurements, blood and urine samples.
- Cancer events and deaths are obtained from the Medical Research Information Service (MRIS) and General Register Office for Scotland (GROS). Links are also being established to hospital data systems.
- A follow-up questionnaire is planned for 2011/12

Health screening design



¹ Point-to-point data transfer via dialup ² overnight ³ Names of participants but no addresses ⁴ via Royal Mail

Samples stored

Type	Content (2 ml / tube)	Cryovials (max.)
EDTA	Plasma	2
	White cells	1
	Red Cells	2
SC	Plasma	1
	White cells	1
	Red Cells	1
SST	Serum	3
	Plasma	2
LHEP	White cells	1
	Red Cells	2
	Urine	5
ACD	ACD	2
Total		23




EDTA = Ethylenediaminetetraacetic acid; SC=Sodium Citrate; SST=Serum SeparatingTube; LHEP=Lithium Heparin; ACD = acid-citrate-dextrose


Progress and Results

- 29 forces have been included to date (17 forces complete; 5 forces – screening ongoing; 7 forces – enrolment phase ongoing), with 27,631 participants screened
- 23 x 27,631 = 635,513 x 2ml aliquots, most stored in liquid-nitrogen (vapour-phase) at Hammersmith Biorepository (pictured above, London)
- Backups in Northwick Park Hospital (London) and Queen Mary, University of London

Number of participants screened per police force by the end of 2010



Percentages of screened participants for various health indicators by age in the Airwave Health Monitoring Study (2010)



DBP Diastolic Blood Pressure; SBP Systolic Blood Pressure; BMI Body Mass Index; WC Waist Circumference; CRP High Sensitivity C-reactive Protein; HbA1C Glycosylated Haemoglobin; ECG Electrocardiogram